arXiv:hep-ph/9612312v1 10 Dec 1996

Yaroslavl State University
Preprint YARU-HE-96/05

hep-ph/9612312

The neutrino energy and momentum loss through the process $\nu \to \nu e^- e^+$ in a strong magnetic field

A.V. Kuznetsov[*], N.V. Mikheev[†]

Division of Theoretical Physics, Department of Physics,
Yaroslavl State University, Sovietskaya 14,
150000 Yaroslavl, Russian Federation.

10 December 1996

Abstract

A process of the electron–positron pair production by neutrino propagating in a strong magnetic field is investigated in the framework of the Standard Model. The process probability and the mean values of the neutrino energy and momentum loss are calculated. Possible astrophysical manifestations of the process considered are briefly analysed.

Submitted to Physics Letters B

---

[*] E-mail address:avkuzn@univ.uniyar.ac.ru
[†] E-mail address: mikheev@yars.free.net

The neutrino interaction with an intensive electromagnetic field plays an important role in neutrino physics. The interaction was generally considered beyond the Standard Model, for example, due to an unusually large neutrino magnetic moment [1]. In the recent times an interesting effect of significant enhancement of the probability of the neutrino radiative decay $\nu_i \to \nu_j \gamma$ by a magnetic field (magnetic catalysis) was discovered in the frame of the Standard Model with lepton mixing [2].

Another decay channel also exists, $\nu_i \to \nu_j e^- e^+$ (i and j enumerate the neutrino mass eigenstates but not the neutrinos with definite flavors), which is forbidden in vacuum when $m_i < m_j + 2m_e$. However, the kinematics of a charged particle in a magnetic field is that which allows to have a sufficiently large space–like total momentum for the electron–positron pair, and this process is possible even for very light neutrinos. It means that a flavor of the ultrarelativistic neutrino is conserved in this transition in a magnetic field, to the terms of the order of m_ν^2/E_ν^2 regardless of the lepton mixing angles. Consequently, a question of neutrino mixing is not pertinent in this case and the process $\nu \to \nu e^- e^+$ can be considered in the frame of the Standard Model without lepton mixing.

In this paper we investigate the process $\nu \to \nu e^- e^+$ for the high energy neutrino, $E_\nu \gg m_e$, in a strong constant magnetic field. We consider a magnetic field as the strong one if it is much greater than the known Schwinger value $B_e = m_e^2/e \simeq 4.41 \cdot 10^{13} G$. This process could be of importance in astrophysical applications, e.g. in an analysis of cataclysms like a supernova explosion or a coalescence of neutron stars, where the strong magnetic fields can exist, and where neutrino processes play the central physical role. In the present view, the magnetic field strength inside the astrophysical objects in principle could be as high as $10^{16} - 10^{18} G$, both for toroidal [3, 4] and for poloidal [5] fields. The magnetic fields of the order of $10^{12} - 10^{13}\, G$ have been observed at the surface of pulsars are the so-called 'old' magnetic fields and they do not provide a determination of the field strength at a moment of the cataclysm [4]. Curiously, a model was proposed recently for soft gamma-ray repeaters to be associated with young neutron stars, with surface dipole fields $B \sim 10^{14} - 10^{15} G$ [6].

Here we calculate the probability of the process and the mean values of the neutrino energy and momentum loss through the production of electron–positron pairs.

If the momentum transferred is relatively small, $|q^2| \ll m_W^2$ [1], the weak interaction of neutrinos with electrons could be described in the local limit by the effective Lagrangian of the form

$$\mathcal{L} \,=\, \frac{G_F}{\sqrt{2}} [\bar{e}\gamma_\alpha (g_V - g_A \gamma_5) e] \, [\bar{\nu}\gamma^\alpha (1-\gamma_5)\nu] \,, \tag{1}$$

where

$$g_V = \pm\frac{1}{2} + 2 sin^2\theta_W \,, \quad g_A = \pm\frac{1}{2}. \tag{2}$$

Here the upper signs correspond to the electron neutrino ($\nu = \nu_e$) when both neutral and charged current interaction takes part in a process. The lower signs correspond to μ and τ neutrinos ($\nu = \nu_\mu, \nu_\tau$), when the neutral current interaction is only presented in the Lagrangian (1).

An amplitude of the process $\nu \to \nu e^- e^+$ could be immediately obtained from the Lagrangian (1) where the known solutions of the Dirac equation in a magnetic field should be used. The general expression for the probability takes a rather complicated form. We present here the results of our calculations in two limiting cases which have a clear physical meaning: $eB > E^2$ and $eB \ll E^2$.

In the case when the field strength B appears to be the largest physical parameter, the electron and the positron could be born only in the states corresponding to the lowest Landau level. Being integrated over the $e^- e^+$ pair phase space with the fixed total pair momentum q, the process probability could be written in the form

$$wE \,=\, \frac{G_F^2 m^2 eB}{32\pi^4} \int \frac{d^3k}{\omega} \, \frac{e^{-q_\perp^2/2eB}}{(q_\parallel^2)^{3/2}(q_\parallel^2 - 4m^2)^{1/2}} \, \left| g_V (q\tilde{\varphi}j) - g_A (q\varphi\varphi j) \right|^2 \,. \tag{3}$$

Hereafter $p^\alpha = (E, \vec{p}),\ k^\alpha = (\omega, \vec{k})$ are the four-momenta of the initial and the final neutrinos, respectively; m is the electron mass; $q = p - k$; $j^\alpha = \bar{\nu}(k)\gamma^\alpha(1-\gamma_5)\nu(p)$; $(q\varphi\varphi j) = q^\alpha \varphi_{\alpha\beta}\varphi^{\beta\delta} j_\delta$, $\varphi_{\alpha\beta}$ is the dimensionless external field tensor, $\varphi_{\alpha\beta} = F_{\alpha\beta}/B$, $\tilde{\varphi}_{\alpha\beta} = \frac{1}{2}\varepsilon_{\alpha\beta\rho\sigma}\varphi^{\rho\sigma}$ is the dual tensor; $q_\parallel^2 = (q\tilde{\varphi}\tilde{\varphi}q)$, $q_\perp^2 = (q\varphi\varphi q)$.

---

[1] As the analysis shows, it corresponds in this case to the neutrino energy $E \ll m_W^3/eB$.

It should be noted that an expression for the probability of the process and in part Eq. (3) could be obtained by another easier way. As is well known, the cross-section of the reaction $\nu\tilde{\nu} \to e^-e^+$ is connected with the imaginary part of the amplitude of the $\nu\tilde{\nu} \to \nu\tilde{\nu}$ transition via the electron loop (see Fig.1) by the unitarity condition

$$\sigma(\nu\tilde{\nu} \to e^-e^+) = \frac{1}{q^2}\, Im\, \mathcal{M}(\nu\tilde{\nu} \to \nu\tilde{\nu}). \tag{4}$$

One can easily see that Eq. (4) allows to find a probability of the crossed channel of the reaction, namely, the process $\nu \to \nu e^-e^+$, by integrating over the phase space of the final neutrino

$$w(\nu \to \nu e^-e^+)E \;=\; \frac{1}{16\pi^3} \int \frac{d^3k}{\omega}\, Im\, \mathcal{M}(\nu\tilde{\nu} \to \nu\tilde{\nu}). \tag{5}$$

Within the above–mentioned limiting case $eB > E^2$ an imaginary part of the amplitude, see Fig.1, corresponds to the virtual electron and positron to become on-shell in the lowest Landau level. A straightforward calculation of Eq. (5) leads to Eq. (3) immediately.

Integrating Eq. (3) over the phase space, one obtains the following expression for the probability

$$\begin{aligned} w(\nu \to \nu e^-e^+) \;&=\; \frac{G_F^2(g_V^2 + g_A^2)}{16\pi^3}\, eBE^3 sin^4\theta \cdot f_1(\lambda), && (6)\\ f_1(\lambda) \;&=\; 1 - \frac{2}{3}\lambda + \frac{5}{16}\lambda^2 - \frac{7}{60}\lambda^3 + \ldots, && (7) \end{aligned}$$

where $\lambda = (p\varphi\varphi p)/eB = E^2 sin^2\theta/eB$, θ is an angle between the initial neutrino momentum $\vec{p}$ and the magnetic field direction.

In another limiting case, $E^2 \gg eB$, when a great number of the Landau levels could be excited, our result can also be represented in a simple form. Using the Eq. (5) we obtain the following expression for the process probability

$$w(\nu \to \nu e^-e^+)E \;=\; \frac{G_F^2(g_V^2 + g_A^2)}{27\pi^3}\, m^6 \chi^2\, (ln\chi - \frac{1}{2} ln3 - \gamma_E - \frac{29}{24}), \tag{8}$$

where $\chi = e(pFFp)^{1/2}/m^3 = \beta E sin\theta/m$ is the so-called field dynamical parameter, $\beta = B/B_e \gg 1$, $\gamma_E = 0.577\dots$ is the Euler constant.

A probability of this process in the last limiting case was obtained earlier in the paper [7] by calculating of an imaginary part of the two-loop amplitude of the $\nu_\mu \to \nu_\mu$ transition via the virtual e^-e^+ pair. However, some numerical errors were committed there. In particular, an extra factor of $\frac{1}{2}$ was inserted. Besides, the calculation technique used in Ref. [7] allows to find an integral probability of the process $\nu \to \nu e^- e^+$ only, but the final neutrino spectrum could not be obtained. Thus, it would not be possible to investigate some interesting astrophysical consequences of this process to be discussed below.

It should be noted that a practical significance of this process for astrophysics could be in the mean values of the neutrino energy and momentum loss rather than in the process probability (6), (8). These mean values could be found from the four-vector

$$Q^\alpha = \frac{1}{16\pi^3} \int \frac{d^3k}{\omega}\, q^\alpha\, Im\, \mathcal{M}(\nu\tilde{\nu} \to \nu\tilde{\nu}). \tag{9}$$

Its zero component is connected with the mean neutrino energy loss in a unit time, $\mathcal{I} = Q_0/E$. The space components of the four-vector (9) are connected similarly with the neutrino momentum loss in unit time, $\vec{\mathcal{F}} = \vec{Q}/E$. Here we present the expressions for Q^α in two limiting cases used above for the probability:

i) $eB \sim E^2$

$$\begin{aligned} Q^\alpha &= \frac{G_F^2 eB (p\varphi\varphi p)^2}{48\pi^3}\{(g_V^2 + g_A^2)[p^\alpha \cdot f_2(\lambda) - 2(\varphi\varphi p)^\alpha \cdot f_3(\lambda)] \\ &+ 2 g_V g_A (\tilde{\varphi} p)^\alpha \cdot f_2(\lambda)\}, \qquad (10) \\ f_2(\lambda) &= 1 - \frac{5}{8}\lambda + \frac{21}{80}\lambda^2 - \frac{7}{80}\lambda^3 + \dots, \\ f_3(\lambda) &= 1 - \frac{15}{16}\lambda + \frac{21}{40}\lambda^2 - \frac{7}{32}\lambda^3 + \dots; \end{aligned}$$

ii) $E^2 \gg eB$

$$Q^\alpha = \frac{7}{16}\frac{wE}{ln\chi - 2.335}\,[\,p^\alpha(ln\chi - 1.888) - \sqrt{3}\,\frac{\beta^2}{\chi}\,(\varphi\varphi p)^\alpha$$

$$- \quad 7.465 \, \frac{g_V g_A}{g_V^2 + g_A^2} \, \frac{\beta}{\chi^{2/3}} \, (\tilde{\varphi} p)^\alpha], \tag{11}$$

where the probability w should be taken from Eq. (8). Eqs. (6), (8), (10), (11) are also applicable for the process with antineutrino $\tilde{\nu} \to \tilde{\nu} e^- e^+$ due to the CP–invariance of the weak interaction.

Let us note that our results are valid in the presence of plasma with the electron density $n \sim 10^{33} - 10^{34} cm^{-3}$, $(n = n_{e^-} - n_{e^+})$. This is due to a peculiarity of the statistics of the relativistic electron gas in a magnetic field [8]. The chemical potential in a very strong field

$$\mu = \frac{2\pi^2 n}{eB} \simeq 0.26 MeV \left(\frac{n}{10^{33} cm^{-3}} \right) \left(\frac{10^{17} G}{B} \right) \tag{12}$$

is significantly less than in the case without field. Because of this, the suppressing statistical factors do not arise in integrating over the phase space of the $e^- e^+$ pair.

To illustrate the formulae obtained we shall consider the astrophysical process of a birth of the magnetized neutron star, pulsar, for example, in a supernova explosion. Let us suppose that in the cataclysm a very strong magnetic field of the order of $10^{16} - 10^{18} G$ [3, 5, 4] arises in any reason in the vicinity of a neutrinosphere. The electron density in this region will be considered to be not too high, so the electron gas could be treated as the nondegenerated one. In this case the neutrino propagating through the magnetic field will loose the energy and the momentum in accordance with our formulas. A part of the total energy lost by neutrinos in the field as well as the total momentum received by a rest of the star from outgoing neutrinos could be estimated from Eq. (10):

$$\frac{\Delta \mathcal{E}}{\mathcal{E}_{tot}} \sim 2 \cdot 10^{-2} \left(\frac{B}{10^{17} G} \right) \left(\frac{\bar{E}}{10 MeV} \right)^3 \left(\frac{\Delta \ell}{10 km} \right), \tag{13}$$

$$\begin{aligned} \vec{\mathcal{P}} &= (0, 0, \mathcal{P}), \\ \mathcal{P} &\sim 3 \cdot 10^{40} \frac{g\, cm}{s} \left(\frac{\mathcal{E}_{tot}}{4 \cdot 10^{53} erg} \right) \left(\frac{B}{10^{17} G} \right) \left(\frac{\bar{E}}{10 MeV} \right)^3 \left(\frac{\Delta \ell}{10 km} \right), \end{aligned} \tag{14}$$

here the z axis is directed along the magnetic field, $\Delta \ell$ is a characteristic size of the region where the field strength varies insignificantly. $\mathcal{E}_{tot}$ is the total energy carried off by neutrinos in a supernova explosion, $\bar{E}$ is the neutrino energy averaged over the neutrino spectrum. Here we take the energy scales which are believed to be typical for supernova explosions [9]. One can see from Eq. (13) that the effect could manifest itself at a level of about percent. In principle, it could be essential in a detailed theoretical description of the process of supernova explosion.

Let us note that an origin of the asymmetry of the neutrino momentum loss with respect to the magnetic field direction is a manifestation of the parity violation in weak interaction, because the $\mathcal{P}$ value is defined by the term in Eq. (10) proportional to the product of the constants g_V and g_A. This asymmetry results in the recoil "kick" velocity of a rest of the cataclysm. If the physical parameters would have in any reason the values of order of the scales in Eq. (14), it would provide a "kick" velocity of order 150 km/s for a pulsar with mass of order of the solar mass.

We are grateful to V.M. Lipunov for helpful discussions, and to A.A. Gvozdev and L.A. Vassilevskaya for useful remarks.

The work of N.V. Mikheev was supported in part by a Grant N d104 from the International Soros Science Education Program.

References

[1] M.B. Voloshin, M.I. Vysotski and L.B. Okun, Zh.Eksp.Teor.Fiz. 91(1986)754 (Sov. Phys. JETP 64(1987)446).

[2] A.A. Gvozdev, N.V. Mikheev and L.A. Vassilevskaya, Phys. Lett. B289(1992)103; B321(1994)108; Yad. Fiz. 57(1994)124 (Phys. At. Nucl. 57(1994)117); Phys. Rev. D54(1996)5674.

[3] M. Ruderman, in "Neutron Stars: Theory and Observation", ed. by J. Ventura and D. Pines, Kluwer Academic. Pub., Dordrecht, 1991.

[4] V.M. Lipunov, Astrophysics of Neutron Stars, Springer-Verlag, New York, 1992, and V.M. Lipunov, private communication.

[5] M. Bocquet et al., Astron. and Astrophys., 301(1995)757.

[6] C. Thompson and R.C. Duncan, Mon.Not.R.Astron.Soc., 275(1995)255.

[7] A.V. Borisov, A.I. Ternov and V.Ch. Zhukovsky, Phys. Lett. B318(1993)489.

[8] L.D. Landau and E.M. Lifshitz, Statistical Physics, Pergamon Press, Oxford, 1980.

[9] V.S. Imshennik and D.K. Nadyozhin, Usp. Fiz. Nauk 156(1988)561 (Sov. Sci. Rev., Sect. E 8(1989)1); D.K. Nadyozhin, in: Particles and Cosmology, Proc. Baksan Int. School, eds. V.A. Matveev et al., World Sci., 1992, p.153.



Fig. 1.

Fig.1. The Feynman diagram for the process $\nu\tilde{\nu} \to \nu\tilde{\nu}$. The double line corresponds to the exact propagator of an electron in a magnetic field.